



A note on a cycle partition problem[☆]

Fengli Yang, Elkin Vumar^{*}

College of Mathematics and System Sciences, Xinjiang University, Urumqi 830046, PR China

ARTICLE INFO

Article history:

Received 3 June 2009

Received in revised form 31 January 2011

Accepted 3 February 2011

Keywords:

Circumference

Partition

Longest cycle

Dominating cycle

Path partition conjecture

ABSTRACT

Let G be any graph, and let $c(G)$ denote the circumference of G . If, for every pair c_1, c_2 of positive integers satisfying $c_1 + c_2 = c(G)$, the vertex set of G admits a partition into two sets V_1 and V_2 such that V_i induces a graph of circumference at most c_i , $i = 1, 2$, then G is said to be c -partitionable. In [M.H. Nielsen, On a cycle partition problem, Discrete Math. 308 (2008) 6339–6347], it is conjectured that every graph is c -partitionable. In this paper, we verify this conjecture for a graph with a longest cycle that is a dominating cycle. Moreover, we prove that G is c -partitionable if $c(G) \geq |V(G)| - 3$.

© 2011 Elsevier Ltd. All rights reserved.

1. Introduction

Throughout this paper, we consider only finite undirected graphs without loops and multiple edges. For notation and terminology not defined here, we refer to West [1].

The vertex set and edge set of a graph G are denoted by $V = V(G)$ and $E = E(G)$, respectively. $\alpha(G)$ and $\kappa(G)$ denote the independence number and the connectivity of G , respectively. If $k \leq \alpha(G)$, we define $\sigma_k(G)$ as the minimum value of the degree sum of any k independent vertices of G ; otherwise, we set $\sigma_k(G) = +\infty$. For $S \subseteq V$, let $\langle S \rangle$ denote the subgraph induced by S . For subgraphs H and K of G , let $G - H := \langle V \setminus V(H) \rangle$, and let $N_H(K)$ denote the set of vertices in H that are adjacent to some vertex in K . Moreover, we let $N(K) := N_{G-K}(K)$. In particular, if K consists of one vertex v , we omit the brackets, and we use $d_H(v) = |N_H(v)|$ and $d(v) = |N(v)|$. A cycle C is called a *dominating cycle* if $G - C$ is edgeless.

The *circumference* $c(G)$ of G is defined as follows. If G is edgeless, then $c(G) = 1$; if G is acyclic but contains an edge, then $c(G) = 2$; finally, if G contains a cycle, then $c(G)$ is the length of a longest cycle in G . A longest path in a graph G is called a *detour* of G . The number of vertices in a detour of G is called the *detour order* of G , and is denoted by $\tau(G)$.

A graph G is said to be τ -partitionable if, for every pair (a, b) of positive integers with $a + b = \tau(G)$, V has a partition $V = V_1 \cup V_2$ such that $\tau(\langle V_1 \rangle) \leq a$ and $\tau(\langle V_2 \rangle) \leq b$.

The following conjecture, known as the Path Partition Conjecture (PPC), was posed by Laborde et al. [2] in 1982.

Conjecture 1. Every graph is τ -partitionable.

In recent years, a number of results have been reported in support of the conjecture (see, e.g., [3–7]). However, the general conjecture seems to be quite difficult to settle. Bondy [8] considered the directed version of the conjecture, and some papers have dealt with that conjecture (see, e.g., [9,10]).

Notice that similar concepts have also been defined for other parameters. A graph G is called Δ -partitionable (where Δ stands for the maximum degree of G) if, for every pair (a, b) of positive integers with $a + b = \Delta - 1$, there is a partition (V_1, V_2)

[☆] This work is supported by NSFC (No. 11061034) and XJEDU2010101.

^{*} Corresponding author.

E-mail address: vumar@xju.edu.cn (E. Vumar).

of V such that $\Delta(\langle V_1 \rangle) \leq a$ and $\Delta(\langle V_2 \rangle) \leq b$. In [11], Lovász proved that every graph is Δ -partitionable. Stiebitz proved a dual-type result with respect to the minimum degree [12]. Recently, Nielsen [13] defined a cycle analog of the path partition problem in a natural way. If c_1 and c_2 are positive integers and (V_1, V_2) is a partition of $V(G)$ such that $c(\langle V_i \rangle) \leq c_i$, $i = 1, 2$, then we say that (V_1, V_2) is a (c_1, c_2) -partition of G and that G is (c_1, c_2) -partitionable. If G is (c_1, c_2) -partitionable for every pair (c_1, c_2) of positive integers with $c_1 + c_2 = c(G)$, then G is said to be c -partitionable. For more details on the PPC and related problems, we refer the reader to [14,3,13] and the references therein.

The following Conjecture 2 was formulated by Nielsen [13].

Conjecture 2 ([13]). Every graph is c -partitionable.

Among other results, the following Theorem 1.1 and Proposition 1.2 have been proved in [13].

Theorem 1.1 ([13]). A graph G is c -partitionable if G belongs to one of the following graph classes.

(a) Planar graphs. (b) Bipartite graphs. (c) Comparability graphs. (d) Chordal graphs. (e) Claw-free graphs. (f) Complete multipartite graphs.

Proposition 1.2 ([13]). Every graph G of circumference at least $|V(G)| - 1$ is c -partitionable.

The following result of Nielsen shows that a graph with circumference at most 9 is also c -partitionable.

Theorem 1.3 ([13]). Let G be a graph, and let $1 \leq c_1 \leq c_2$ be positive integers such that $c_1 + c_2 = c(G)$ and $c_1 \leq 4$. Then G is (c_1, c_2) -partitionable.

In this paper, we prove that a graph is c -partitionable if it admits a longest cycle which is a dominating cycle.

Theorem 1.4. If a graph G admits a longest cycle that is a dominating cycle, then G is c -partitionable.

Moreover, we extend Proposition 1.2 by decreasing the upper bound on the circumference of a c -partitionable graph to $|V(G)| - 3$. Note that Frick and Schiermeyer proved that a graph G with detour order at least $|V(G)| - 3$ is τ -partitionable [6].

Proposition 1.5. Every graph G of circumference at least $|V(G)| - 3$ is c -partitionable.

There are some Dirac-type and Ore-type conditions for a graph to admit a dominating longest cycle (see [15–22]). By Theorem 1.4, all these are the conditions for a graph to be c -partitionable, and hence we have the following two corollaries.

Corollary 1.6. Let G be a 2-connected graph of order n . G is c -partitionable if one of the following holds.

(a) $\delta \geq \frac{n-3}{2}$. (b) $\delta \geq \frac{n+2}{3}$. (c) $\sigma_2(G) \geq n - 3$. (d) $\sigma_3(G) \geq n + 2$.

Corollary 1.7. Let G be a 3-connected graph of order n . G is c -partitionable if one of the following holds.

(a) $\sigma_4 \geq n + 2\kappa$. (b) $n \geq 13$, $\sigma_4(G) \geq \frac{4}{3}n + \frac{5}{3}$. (c) $\sigma_4(G) \geq \frac{3}{2}n + 1$. (d) $\sigma_4(G) \geq n + \kappa + 3$.

2. Proof of the main results

Proof of Theorem 1.4. If $c(G) \leq 2$, then G is bipartite, and, by Theorem 1.1, G is c -partitionable. So we may assume that $c(G) \geq 3$ and that G has a longest cycle that is a dominating cycle. Then, obviously, it suffices to consider the case when G is connected.

Suppose that G is a connected graph and that C is a longest cycle (with a given cyclic orientation) that is a dominating cycle. Then $G - C = \{v_1, v_2, \dots, v_k\}$ is an independent set. Suppose that $1 \leq c_1 \leq c_2$ are two integers with $c_1 + c_2 = c(G)$. For distinct vertices x, y on C , we use $C(x, y)$ to denote the set of vertices on C from x to y (not including x and y) with respect to the orientation. From the maximality of C , it is not difficult to get the following observations.

Observation. (a) If $x, y \in N(v_i)$ with $x \neq y$, then $|C(x, y)| \geq 1$.

(b) $|N(v_i)| \leq \left\lfloor \frac{c(G)}{2} \right\rfloor$, $i = 1, 2, \dots, k$.

(c) Let $x_1, x_2 \in N(v_i), x'_1, x'_2 \in N(v_j)$, $i \neq j$, and let these four distinct vertices appear on C in the following order: x_1, x'_1, x_2, x'_2 . Then $|C(x_1, x'_1) \cup C(x_2, x'_2)| \geq 2$ and $|C(x'_1, x_2) \cup C(x'_2, x_1)| \geq 2$.

Case 1. $\left| \bigcup_{i=1}^k N(v_i) \right| \leq c_2$.

In this case, since $\left| \bigcup_{i=1}^k N(v_i) \right| \leq c_2$, we let V_2 be a set of order c_2 satisfying $\bigcup_{i=1}^k N(v_i) \subseteq V_2 \subseteq C$, and let $V_1 = V(G) \setminus V_2$.

Obviously, $c(\langle V_2 \rangle) \leq |V_2| = c_2$. Since every vertex of $G - C$ is of degree zero in $\langle V_1 \rangle$, v_i is not on any cycle in $\langle V_1 \rangle$, for $i = 1, 2, \dots, k$. Hence, we have $c(\langle V_1 \rangle) \leq |V_1 - \{v_1, \dots, v_k\}| = c_1$. Thus (V_1, V_2) is a (c_1, c_2) -partition of G .

Case 2. $\left| \bigcup_{i=1}^k N(v_i) \right| > c_2$.

Claim 2.1. $c(G) \geq 2 \left| \bigcup_{i=1}^k N(v_i) \right| - k$.

Proof of Claim 2.1. First, suppose that $N(v_i) \cap N(v_j) = \emptyset$ for $i \neq j$. By observation, it is easy to see that

$$\begin{aligned} c(G) &\geq |N(v_1)| + |N(v_2)| + \cdots + |N(v_k)| + (|N(v_1)| - 1) + \cdots + (|N(v_k)| - 1) \\ &= 2 \left| \bigcup_{i=1}^k N(v_i) \right| - k. \end{aligned}$$

Next, suppose that there are distinct vertices $v_i, v_j \in G - C$ such that $N(v_i) \cap N(v_j) \neq \emptyset$. Set $N'(v_1) = N(v_1)$, $N'(v_2) = N(v_2) \setminus N(v_1)$, \dots , $N'(v_k) = N(v_k) \setminus \bigcup_{i=1}^{k-1} N(v_i)$. Then, again by observation, we have

$$\begin{aligned} c(G) &\geq |N'(v_1)| + |N'(v_2)| + \cdots + |N'(v_k)| + (|N'(v_1)| - 1) + \cdots + (|N'(v_k)| - 1) \\ &= 2(|N'(v_1)| + |N'(v_2)| + \cdots + |N'(v_k)|) - k \\ &= 2 \left| \bigcup_{i=1}^k N(v_i) \right| - k. \quad \square \end{aligned}$$

Claim 2.2. $\left| \bigcup_{i=1}^k N(v_i) \right| \leq \left\lceil \frac{c(G)}{2} \right\rceil + \left\lfloor \frac{k}{2} \right\rfloor$.

Proof of Claim 2.2. From Claim 2.1, we know that $\left| \bigcup_{i=1}^k N(v_i) \right| \leq \frac{c(G)+k}{2}$. Since $\left| \bigcup_{i=1}^k N(v_i) \right|$ is an integer, we have $\left| \bigcup_{i=1}^k N(v_i) \right| \leq \left\lfloor \frac{c(G)+k}{2} \right\rfloor \leq \left\lceil \frac{c(G)}{2} \right\rceil + \left\lfloor \frac{k}{2} \right\rfloor$. \square

Since $\left\lceil \frac{c(G)}{2} \right\rceil \leq c_2$, by Claim 2.2, we have $\left| \bigcup_{i=1}^k N(v_i) \right| \leq c_2 + \left\lfloor \frac{k}{2} \right\rfloor$. Without loss of generality, we may assume that $d_G(v_1) \geq d_G(v_2) \geq \cdots \geq d_G(v_k)$. We choose a vertex $u_1 \in N(v_1)$ such that $d_{G-C}(u_1) = \max\{d_{G-C}(w) | w \in N(v_1)\}$, and let $u_1 \in V_2$. Now, consider the graph $G_1 = G - u_1$. In G_1 , we relabel v_1, v_2, \dots, v_k as v'_1, v'_2, \dots, v'_k such that $d_{G_1}(v'_1) \geq d_{G_1}(v'_2) \geq \cdots \geq d_{G_1}(v'_k)$. Then, we choose $u_2 \in N_{G_1}(v'_1)$ such that $d_{G_1-C}(u_2) = \max\{d_{G_1-C}(w) | w \in N_{G_1}(v'_1)\}$ and let $u_2 \in V_2$. Continue this process until $|V_2| = c_2$. Note this is possible because $\left| \bigcup_{i=1}^k N(v_i) \right| > c_2$. Obviously, $V_2 \subseteq \bigcup_{i=1}^k N(v_i)$. Further, since $c_2 < \left| \bigcup_{i=1}^k N(v_i) \right| \leq c_2 + \left\lfloor \frac{k}{2} \right\rfloor$, we know that $\left| \bigcup_{i=1}^k N(v_i) - V_2 \right| \leq \left\lfloor \frac{k}{2} \right\rfloor$. Let $V_1 = V \setminus V_2$. Then, by Claim 2.2 and the choice of V_2 , it is not difficult to see that each v_i has degree at most 1 in $\langle V_1 \rangle$, and consequently $c(\langle V_1 \rangle) \leq |V_1 - \{v_1, v_2, \dots, v_k\}| = c_1$. So G is (c_1, c_2) -partitionable.

The proof of Theorem 1.4 is now complete. \square

Proof of Proposition 1.5. By Proposition 1.2, it suffices to prove the cases $c(G) = |V(G)| - 2$ and $c(G) = |V(G)| - 3$. We only prove the case $c(G) = |V(G)| - 3$, since the other case is similar (and even simpler). Let G be a graph with $c(G) = c_1 + c_2$, $1 \leq c_1 \leq c_2$, and let C be a longest cycle in G with $|V(G - C)| = 3$. In view of Theorem 1.3, we may assume that $c_1 \geq 5$.

Using similar arguments as in the proofs of Claims 2.1 and 2.2, one can obtain $|N_C(G - C)| \leq c_2 + 1$. If the strict inequality holds, then, using similar arguments as those in Case 1 of the proof of Theorem 1.4, we can show that G has a (c_1, c_2) -partition. So, we may assume that $|N_C(G - C)| = c_2 + 1$. We let V_2 be any set of order c_2 satisfying $|N_C(G - C) \setminus V_2| = 1$, and let $V_1 = V \setminus V_2$. Then, from the choice of V_2 , it follows that $|N_C(G - C) \cap V_1| = 1$, and this in turn implies that in $\langle V_1 \rangle$ the three vertices of $G - C$ cannot be on any cycle of length greater than 4. Since $c_1 \geq 5$, we have $c(\langle V_1 \rangle) \leq \max\{c(\langle V_1 \setminus V(G - C) \rangle), 4\} \leq |V_1 \setminus V(G - C)| = c_1$. The proposition is proved. \square

Acknowledgement

The authors would like to thank the reviewer for his/her valuable comments and suggestions, which have greatly improved both the exposition and the clarity of this paper.

References

- [1] D.B. West, Introduction to Graph Theory, second ed., Prentice Hall, Upper Saddle River, NJ, 2001.
- [2] J.M. Laborde, C. Payan, N.H. Xuong, Independent sets and longest directed paths in digraphs, in: Graphs and Other Combinatorial Topics, Prague, 1982, in: Teubner-Texte Math., vol. 59, 1983, pp. 173–177.
- [3] J.E. Dunbar, M. Frick, The path partition conjecture is true for claw-free graphs, Discrete Math. 307 (11–12) (2007) 1285–1290.
- [4] J. Dunbar, M. Frick, Path kernels and partitions, J. Combin. Math. Combin. Comput. 31 (1999) 137–149.
- [5] J.E. Dunbar, M. Frick, F. Bullock, Path partitions and P_n -free sets, Discrete Math. 289 (1–3) (2004) 145–155.
- [6] M. Frick, I. Schiermeyer, An asymptotic result for the path partition conjecture, Electron. J. Combin. 10 (2005) Research Paper 48 (electronic).
- [7] M. Frick, C. Whitehead, A new approach for the path partition conjecture, Util. Math. 69 (2006) 195–206.
- [8] J.A. Bondy, in: R.L. Graham, M. Grötschel, L. Lovász (Eds.), Handbook of Combinatorics, vol. I, The MIT Press, Cambridge, MA, 1995, p. 49.

- [9] S. van Aardt, G. Dlamini, J. Dunbar, M. Frick, O. Oellermann, The directed path partition conjecture, *Discuss. Math. Graph Theory* 25 (2005) 331–343.
- [10] J. Bang-Jensen, M.H. Nielsen, A. Yeo, Longest path partitions in generalizations of tournaments, *Discrete Math.* 306 (2006) 1830–1839.
- [11] L. Lovász, On decomposition of graphs, *Studia Sci. Math. Hungar.* 1 (1966) 237–238.
- [12] M. Stiebitz, Decomposing graphs under degree constraints, *J. Graph Theory* 23 (1996) 321–324.
- [13] M.H. Nielsen, On a cycle partition problem, *Discrete Math.* 308 (2008) 6339–6347.
- [14] R.E.L. Aldred, C. Thomassen, Graphs with not all possible path-kernels, *Discrete Math.* 285 (2004) 297–300.
- [15] J.C. Bermond, On Hamiltonian walks, *Congr. Numer.* 15 (1976) 41–51.
- [16] G.A. Dirac, Some theorems on abstract graphs, *Proc. Lond. Math. Soc.* 2 (1952) 69–81.
- [17] H. Enomoto, J. van den Heuvel, A. Kaneko, A. Saito, Relative length of long paths and cycles in graphs with large degree sums, *J. Graph Theory* 20 (1995) 213–225.
- [18] R. Li, A. Saito, R.H. Schelp, Relative length of longest paths and cycles in 3-connected graphs, *J. Graph Theory* 37 (2001) 137–156.
- [19] M. Lu, H. Liu, F. Tian, Relative length of longest paths and cycles in graphs, *Graphs Combin.* 23 (2007) 433–443.
- [20] C.St.J.A. Nash-Williams, Edge-disjoint Hamiltonian circuits in graphs with vertices of large valency, in: *Studies in Pure Mathematics*, Academic Press, London, 1971, pp. 157–183.
- [21] Z. Sun, F. Tian, B. Wei, Degree sums, connectivity and dominating cycles in graphs, *Graphs Combin.* 17 (2001) 555–564.
- [22] T. Yamashita, Degree sum and connectivity conditions for dominating cycles, *Discrete Math.* 308 (2008) 1620–1627.